

# University of Georgia Herbicide Programs for Tropical Spiderwort (Benghal Dayflower) Control in 2013 Cotton

A. S. Culpepper, J. T. Flanders, T. M. Webster


Tropical spiderwort is a noxious, exotic, invasive weed that can spread quickly. Upon initial observation, tropical spiderwort appears to be a grass (Figure 1). While not a grass, it is a monocot (in contrast to broadleaf weeds, which are dicots) with leaves and stems usually fleshy and succulent. The stems will creep along the ground and root at the nodes. Vegetative cuttings from stems are capable of rooting and reestablishing following cultivation. Tropical spiderwort will produce seed above and below ground (Figure 2).

Prior to glyphosate-resistant Palmer amaranth challenging our cotton industry, this weed was the most problematic weed of cotton. However, Roundup-based programs developed to control Palmer amaranth have indirectly controlled tropical spiderwort very effectively. During 2011/2012 with greater adoption of Liberty-based programs, tropical spiderwort became more problematic as Liberty is not effective in controlling this weed. A programs approach is needed for both Roundup and Liberty based systems. Tables 1 through 3 are suggestions for the management of tropical spiderwort in Georgia cotton. Refer to the most recent cotton production guide or pest control handbook for herbicide rates and proper cotton sizes at time of application.

**Figure 1. Newly emerged.**



**Figure 2. Underground fruiting structures.**



**Table 1. Managing Tropical Spiderwort in Roundup Ready Cotton.<sup>1</sup>**

Preemergence	POST 1	POST 2	Layby Directed
<b>Warrant<sup>2</sup></b> + herbicides <i>appropriate for other weeds</i>	<b>Roundup</b> + <b>Staple, Dual Magnum<sup>3</sup> or Warrant<sup>2</sup></b>  <i>(Use Staple if spiderwort or Palmer amaranth are up)</i>	<b>Roundup</b> + <b>Dual Magnum<sup>3</sup></b>	<b>Direx + MSMA</b>  <i>(Add Aim if spiderwort is greater than 3 inches; the addition of Dual Magnum<sup>3</sup> or Warrant<sup>2</sup> would improve residual control)</i>

<sup>1</sup> Deep turning the land will provide fair control of spiderwort.

<sup>2</sup> Apply Warrant no more than twice per season, once topically. Dual Magnum can be applied topically once per crop.

<sup>3</sup> Dual Magnum can be applied topically once per crop.

**Table 2. Managing Tropical Spiderwort in GlyTol/Liberty Link Cotton.<sup>1</sup>**

Preemergence	POST 1	POST 2	Layby Directed
Warrant <sup>2</sup> + herbicides appropriate for other weeds	<b>Liberty + Staple, Dual Magnum<sup>3</sup> or Warrant<sup>2</sup></b> OR <b>Roundup + Staple, Dual Magnum<sup>3</sup> or Warrant<sup>2</sup></b>	<b>Liberty + Staple, Dual Magnum<sup>3</sup> or Warrant<sup>2</sup></b> OR <b>Roundup + Staple, Dual Magnum<sup>3</sup> or Warrant<sup>2</sup></b>	<b>Direx + MSMA</b>  <i>(Add Aim if spiderwort is greater than 3 inches; the addition of Dual Magnum<sup>3</sup> or Warrant<sup>2</sup> would improve residual control)</i>
	<p><i>Considerations for selecting POST 1 and 2 treatments:</i></p> <p>1. Use Liberty instead of Roundup if Palmer amaranth has emerged.</p> <p>2. With Liberty or Roundup: add Staple if spiderwort has emerged but use Dual or Warrant if spiderwort has not emerged.</p>		

<sup>1</sup> Deep turning the land will provide fair control of spiderwort.

<sup>2</sup> Apply Warrant no more than twice per season, once topically.

<sup>3</sup> Dual Magnum can be applied topically once per crop.

**Table 3. Managing Tropical Spiderwort with Liberty Link Cotton.<sup>1</sup>**

Preemergence	POST 1	POST 2	Layby Directed
Warrant <sup>2</sup> + herbicides appropriate for other weeds	<b>Liberty + Staple</b> <i>(spiderwort emerged)</i>	<b>Liberty + Dual Magnum<sup>3</sup></b>  <i>(before spiderwort not emerges)</i>	<b>Direx + MSMA</b>  <i>(Add Aim if spiderwort is greater than 3 inches; the addition of Dual Magnum<sup>2</sup> or Warrant<sup>3</sup> would improve residual control)</i>
	<b>Liberty + Dual Magnum<sup>3</sup> or Warrant<sup>2</sup></b>  <i>(spiderwort not emerged)</i>		

<sup>1</sup> Deep turning the land will provide fair control of spiderwort.

<sup>2</sup> Apply Warrant no more than twice per season, once topically.

<sup>3</sup> Dual Magnum can be applied topically once per crop.



The University of Georgia and Ft. Valley State University, the U.S. Department of Agriculture and counties of the state cooperating, Cooperative Extension, the University of Georgia College of Agricultural and Environmental Sciences, offers educational programs, assistance and materials to all people without regard to race, color, national origin, age, gender or disability.

**An Equal Opportunity Employer/Affirmative Action Organization**  
**Committed to a Diverse Work Force**